arXiv:math/9905197v1 [math.GT] 31 May 1999

HOMEOMORPHISMS OF ONE-DIMENSIONAL INVERSE LIMITS WITH APPLICATIONS TO SUBSTITUTION TILINGS, UNSTABLE MANIFOLDS, AND TENT MAPS

MARCY BARGE, JAMES JACKLITCH, AND GIOIA VAGO

ABSTRACT. Suppose that f and g are Markov surjections, each defined on a wedge of circles, each fixing the branch point and having the branch point as the only critical value. We show that if the points in the inverse limit spaces associated with f and g corresponding to the branch point are distinguished then these inverse limit spaces are homeomorphic if and only if the substitutions associated with f and g are weakly equivalent. This, and related results, are applied to one-dimensional substitution tiling spaces, one-dimensional unstable manifolds of hyperbolic sets, and inverse limits of tent maps with periodic critical points.

0. INTRODUCTION

Many of the one-dimensional spaces that arise as invariant sets in dynamical systems can be effectively modeled using inverse limits. This approach was first employed by R.F. Williams in a description of one-dimensional hyperbolic attractors [W1]. In that characterization, Williams showed that some power of a diffeomorphism restricted to a one-dimensional hyperbolic attractor is topologically conjugate to the natural (shift) homeomorphism on an inverse limit whose bonding map is an immersion on a wedge of circles. Further [W2], Williams introduced the notion of shift equivalence of maps to classify such natural homeomorphisms up to topological conjugacy. In this paper we address the classification, up to homeomorphism, of inverse limit spaces whose bonding maps act on wedges of circles. A weaker notion than shift equivalence, namely weak equivalence, will arise in this context. As far as we know, the relation of weak equivalence of matrices first arose in [BD]. Consequences of weak equivalence of matrices and its relation to dimension groups were considered in [SV] and the notion has arisen in the classification of almost finite dimensional algebras up to stable isomorphism (see [BJKR], where weak equivalence of matrices, transposed, is called C^*-equivalence). Weak equivalence of matrices will play a part in the theorems we prove here but a stronger equivalence relation, weak equivalence of substitutions, will play the larger role. Our main result (Theorem 1.16) asserts that certain inverse limit spaces are homeomorphic if and only if the associated substitutions are weakly equivalent.

1991 *Mathematics Subject Classification.* 54F15, 54H20, 58F03, 58F12.

The first author was supported in part by NSF-DMS-9627026.

The second author was supported in part by NSF-EPSCoR-93-50-546.

The authors would like to thank Christian Bonatti for fruitful conversations and the Laboratoire de Topologie, Université de Bourgogne, for its warm hospitality.

Our motivation for considering inverse limits of maps on wedges of circles comes from several sources. One is the classification of one-dimensional hyperbolic attractors, up to homeomorphism, as developed in [J]. Others are: one-dimensional substitution tiling spaces; one-dimensional unstable manifolds of hyperbolic basic sets; and a conjecture of Ingram regarding inverse limits of tent maps. These topics will be addressed in the final section of this paper.

1. Main Results

For $n = 1, 2, \dots$, let $X_n = \bigvee_{i=1}^n S_i^1$ denote n topological circles, S_i^1, joined together at a common point, which we denote by b. Assume that each S_i^1 is given an orientation. A map (by which we will always mean a continuous function) $f : X \to Y$ is *monotone* provided $f^{-1}(\{y\})$ is connected for each $y \in Y$ and a map $h : X_n \to X_m$ is said to be *piecewise monotone* provided X_n is a union of finitely many arcs restricted to each of which h is monotone. A piecewise monotone map is *strictly piecewise monotone* if it is not constant on any nontrivial arc. The point $c \in X_n$ is a *critical point* of the map $h : X_n \to X_m$ provided h is not one-to-one in any neighborhood of c and $v \in X_m$ is a *critical value* of h if $v = h(c)$ for some critical point $c \in X_n$.

For $n, m = 1, 2, \dots$, let $\mathcal{M}_{n,m}$ denote the collection of all piecewise monotone maps $f : X_n \to X_m$ that satisfy: $f(b) = b$; b is the only critical value (if any) of f; f is surjective; and f is not constant on any S_i^1, $i = 1, \dots, n$. Let $\mathcal{M} = \bigcup_{n=1}^\infty \mathcal{M}_{n,n}$. We are interested in the topology of the *inverse limit space*, $\varprojlim f$, with *bonding maps* $f \in \mathcal{M}$. More generally, if $f_i : X_n \to X_n$, $i = 1, 2, \dots$, are maps in $\mathcal{M}_{n,n}$, we will denote by $\varprojlim f_i$ the space $\{(x_0, x_1, \dots) \in \prod_{k=0}^\infty X_n : f_i(x_i) = x_{i-1} \text{ for } i = 1, 2, \dots\}$. The topology on $\varprojlim f_i$ is inherited from the product space $\prod_{k=0}^\infty X_n$; with this topology $\varprojlim f_i$ is a *continuum* (compact, connected, metrizable). We will denote by π_k the projection $\pi_k : \varprojlim f_i \to X_n$, $\pi_k(x_0, x_1, \dots, x_k, \dots) = x_k$ and, when all bonding maps are the same, $f_i = f$ for each i, $\hat{f} : \varprojlim f \to \varprojlim f$ will denote the natural (shift) homeomorphism $\hat{f}(x_0, x_1, \dots) = (f(x_0), x_0, x_1, \dots)$.

For each $n = 1, 2, \dots$, let $\mathcal{A}_n$ denote the alphabet consisting of $2n$ letters $\mathcal{A}_n = \{a_1, \dots, a_n, \bar{a}_1, \dots, \bar{a}_n\}$ and let $\mathcal{W}(\mathcal{A}_n)$ be the collection of all finite nonempty words with letters in $\mathcal{A}_n$. Given the k-letter word $b_1 \dots b_k \in \mathcal{W}(\mathcal{A}_m)$, let $\overline{b_1 \dots b_k} = \bar{b}_k \dots \bar{b}_1 \in \mathcal{W}(\mathcal{A}_m)$ with the convention that $\bar{\bar{b}}_i = b_i$. Starting from $f \in \mathcal{M}_{n,m}$, we want to define an associated substitution $\chi_f : \mathcal{A}_n \to \mathcal{W}(\mathcal{A}_m)$ in such a way that $\chi_f(a_i)$ (resp. $\chi_f(\bar{a}_i)$) describes the way in which f maps the circle S_i^1 (resp. the circle S_i^1 oriented in the opposite way) around each S_j^1, keeping track of order and orientation. For this reason, denote by $\dot{J}_1 < \dot{J}_2 < \dots < \dot{J}_k$ the components of $S_i^1 \backslash f^{-1}(\{b\})$, the ordering of such arcs determined by the positive orientation on S_i^1. Then define $\chi_f(a_i) = b_1 \dots b_k$ provided f maps $\dot{J}_j$ homeomorphically onto $\dot{S}_\ell^1 \equiv S_\ell^1 \backslash \{b\}$ in an orientation preserving (resp. reversing) way if and only if $b_j = a_\ell$ (resp. $\bar{a}_\ell$). Recall the meaning of a substitution. Now, when S_i^1 is oriented according to the opposite orientation, $S_i^1 \backslash f^{-1}(\{b\})$ has components $\dot{J}_k < \dot{J}_{k-1} < \dots < \dot{J}_1$. Then it is natural to define $\chi_f(\bar{a}_i) \equiv \bar{b}_k \dots \bar{b}_1 = \overline{\chi_f(a_i)}$. We then extend χ_f to $\chi_f : \mathcal{W}(\mathcal{A}_n) \to \mathcal{W}(\mathcal{A}_m)$ by concatenation: $\chi_f(c_1 \dots c_\ell) = \chi_f(c_1) \dots \chi_f(c_\ell)$. Note that $\chi_f(\overline{c_1 \dots c_\ell}) = \overline{\chi_f(c_1 \dots c_\ell)}$. For any substitution $\tau : \mathcal{A}_n \to \mathcal{W}(\mathcal{A}_m)$ satisfying $\tau(\bar{b}) = \overline{\tau(b)}$, let A_τ be the $m \times n$ matrix whose ji-th entry is $(A_\tau)_{ji} = \#\{\ell : b_\ell \in \{a_j, \bar{a}_j\} \text{ where } \tau(a_i) = b_1 \dots b_k\}$. That is, $(A_\tau)_{ji}$ is the number of

occurrences of the letters a_j and $\bar{a}_j$ in the word $\tau(a_i)$. In case $\tau = \chi_f$, we will say that f *follows the pattern* τ and we will write A_f for A_τ.

Suppose that $\chi : \mathcal{A}_n \to \mathcal{W}(\mathcal{A}_n)$ and $\psi : \mathcal{A}_m \to \mathcal{W}(\mathcal{A}_m)$ are substitutions that satisfy $\chi(\bar{b}) = \overline{\chi(b)}$ and $\psi(\bar{c}) = \overline{\psi(c)}$ for all $b \in \mathcal{A}_n$, $c \in \mathcal{A}_m$. We will say that χ and ψ are *weakly equivalent*, $\chi \sim_w \psi$, if there are sequences of positive integers $\{n_i\}$, $\{m_i\}$, and substitutions $\sigma_i : \mathcal{A}_m \to \mathcal{W}(\mathcal{A}_n)$, $\tau_i : \mathcal{A}_n \to \mathcal{W}(\mathcal{A}_m)$, $i = 1, 2, \dots$, such that $\sigma_i(\bar{c}) = \overline{\sigma_i(c)}$, $\tau_i(\bar{b}) = \overline{\tau_i(b)}$ for all $c \in \mathcal{A}_m$, $b \in \mathcal{A}_n$ and, after extending the substitutions by concatenation, $\sigma_i \circ \tau_i = \chi^{n_i}$, $\tau_i \circ \sigma_{i+1} = \psi^{m_i}$. That is, the following infinite diagram commutes.



If $A_{n\times n}$ and $B_{m\times m}$ are square matrices with nonnegative integer entries, we will say that A and B are *weakly equivalent*, $A \sim_w B$, provided there are sequences of positive integers $\{n_i\}$, $\{m_i\}$, and nonnegative integer matrices S_i, T_i, $i = 1, 2, \dots$, such that $S_i T_i = A^{n_i}$ and $T_i S_{i+1} = B^{m_i}$. (Weak equivalence of A and B is called C^*-equivalence of A transpose and B transpose in [BJKR].)

Lemma 1.1. *If $f, g \in \mathcal{M}$ and $\chi_f \sim_w \chi_g$ then $A_f \sim_w A_g$.*

Proof. If n_i, m_i, σ_i, τ_i realize the weak equivalence of χ_f and χ_g, let S_i and T_i be given by $S_i = A_{\sigma_i}$ and $T_i = A_{\tau_i}$. It is immediate that $A_f^{n_i} = S_i T_i$ and $A_g^{m_i} = T_i S_{i+1}$. □

Lemma 1.2. *Given $f \in \mathcal{M}_{n,n}$ there are strictly piecewise monotone $f_i \in \mathcal{M}_{n,n}$, $i = 1, 2, \dots$, each of which follows the pattern of χ_f, such that $\varprojlim f_i \simeq \varprojlim f$.*

Proof. The approximation theorem of Mort Brown ([Bro]) asserts that there are $\epsilon_i > 0$ so that if $f_i : X_n \to X_n$ is uniformly within ϵ_i of f (fix some metric on X_n) for each i then $\varprojlim f_i \simeq \varprojlim f$. Clearly, such f_i can be chosen to be strictly piecewise monotone, in $\mathcal{M}_{n,n}$, and follow the pattern of χ_f. □

Lemma 1.3. *If $f, g \in \mathcal{M}_{n,n}$ and $\chi_f = \chi_g$ then $\varprojlim f \simeq \varprojlim g$.*

Proof. Let $f_i, g_i \in \mathcal{M}_{n,n}$ be as in Lemma 1.2 with $\varprojlim f_i \simeq \varprojlim f$ and $\varprojlim g_i \simeq \varprojlim g$, all following the pattern of $\chi_f = \chi_g$. Let $P_0 : X_n \to X_n$ be the identity. Suppose that, for $j = 0, \dots, k$, homeomorphisms $P_j : X_n \to X_n$ have been defined in such a way that $P_j(S_i^1) = S_i^1$ for each $i \in \{1, \dots, n\}$ and the diagram



commutes. Since $f_{k+1}, g_{k+1} \in \mathcal{M}_{n,n}$ are strictly piecewise monotone, and they follow the same pattern, for each i, $S^1_i \backslash f^{-1}_{k+1}(\{b\}) = \bigcup_{\ell=1}^{r_i} \dot{J}_{i,\ell}$ and $S^1_i \backslash g^{-1}_{k+1}(\{b\}) = \bigcup_{\ell=1}^{r_i} \dot{J}'_{i,\ell}$ where the $\dot{J}_{i,\ell}$ and $\dot{J}'_{i,\ell}$ are pairwise disjoint open arcs, $\bigcup_{\ell=1}^{r_i} \dot{J}_{i,\ell}$ and $\bigcup_{\ell=1}^{r_i} \dot{J}'_{i,\ell}$ are dense in S^1_i, $\dot{J}_{i,1} < \dot{J}_{i,2} < \ldots < \dot{J}_{i,r_i}$ and $\dot{J}'_{i,1} < \dot{J}'_{i,2} < \ldots < \dot{J}'_{i,r_i}$ (the order given by the positive orientation on S^1_i), $f_{k+1}|_{\dot{J}_{i,\ell}}$ and $g_{k+1}|_{\dot{J}'_{i,\ell}}$ are both orientation preserving or both orientation reversing homeomorphisms, and $f_{k+1}(\dot{J}_{i,\ell}) = \dot{S}^1_r = g_{k+1}(\dot{J}'_{i,\ell})$ for some $r = r(i,\ell)$. Let $P_{k+1}|_{\dot{J}_{i,\ell}} : \dot{J}_{i,\ell} \to \dot{J}'_{i,\ell}$ be defined by $P_{k+1}|_{\dot{J}_{i,\ell}} = (g_{k+1}|_{\dot{J}'_{i,\ell}})^{-1} \circ P_k \circ f_{k+1}|_{\dot{J}_{i,\ell}}$. P_{k+1} then extends from $\bigcup_{i=1}^{n} \bigcup_{j=1}^{r_i} \dot{J}_{i,j}$ to a homeomorphism of X_n onto itself so that $P_k \circ f_{k+1} = g_{k+1} \circ P_{k+1}$.



The preceding commuting diagram then induces a homeomorphism $\hat{P} : \varprojlim f_i \to \varprojlim g_i$ by $\hat{P}((x_0, x_1, \ldots)) = (P_0(x_0), P_1(x_1), \ldots)$. □

Lemma 1.4. *Given $f \in \mathcal{M}_{n,n}$ there is a strictly piecewise monotone $f' \in \mathcal{M}_{n,n}$ that follows the same pattern as f with $\varprojlim f' \simeq \varprojlim f$.*

Proof. This follows immediately from Lemma 1.3 by taking, for instance, f' to be any of the f_i of 1.2. □

Theorem 1.5. *If $f, g \in \mathcal{M}$ with $\chi_f \sim_w \chi_g$ then $\varprojlim f \simeq \varprojlim g$.*

Proof. By Lemma 1.4, we may assume that f and g are strictly piecewise monotone. Suppose that $f \in \mathcal{M}_{n,n}$, $g \in \mathcal{M}_{m,m}$ and let $\{n_i\}$, $\{m_i\}$, $\sigma_i : \mathcal{A}_m \to \mathcal{W}(\mathcal{A}_n)$, $\tau_i : \mathcal{A}_n \to \mathcal{W}(\mathcal{A}_m)$ be such that, after extending the substitutions by concatenation, $\sigma_i \tau_i = \chi_f^{n_i}$, $\tau_i \sigma_{i+1} = \chi_g^{m_i}$. We will construct continuous surjections $s_i : X_m \to X_n$ and $t_i : X_n \to X_m$ such that $s_i \circ t_i = f^{n_i}$ and $t_i \circ s_{i+1} = g^{m_i}$.

Let $s_1 : X_m \to X_n$ be any continuous strictly piecewise monotone surjection that takes b to b and follows the pattern σ_1. Suppose that $s_i : X_m \to X_n$, $i = 1, \ldots, N$, and $t_i : X_n \to X_m$, $i = 1, \ldots, N-1$, are all continuous strictly piecewise monotone surjections taking b to b following the patterns σ_i and τ_i that have been defined so that the following diagram commutes.



Now $(f^{n_N})^{-1}(\{b\})$ breaks each $S^1_i \subset X_n$ into a certain number of intervals; we define the same number of intervals in X_m in two steps, with the aid of the substitutions σ_N and τ_N. Fix $i \in \{1, \ldots, n\}$ and suppose that $\chi_f^{n_N}(a_i) = b_1 \ldots b_s$.

Then $(f^{n_N})^{-1}(\{b\}) \cap S^1_i = \{b = x_0 < x_1 < \ldots < x_s = b\}$ with f^{n_N} mapping (x_{j-1}, x_j) homeomorphically onto $\dot{S}^1_\ell$ in an orientation preserving (resp. reversing) way iff $b_j = a_\ell$ (resp. $\bar{a}_\ell$). Let $\tau_N(a_i) = c_1 \ldots c_k$. Since $\sigma_N \tau_N(a_i) = \chi_f^{n_N}(a_i) = b_1 \ldots b_s$, there are integers $0 = p_0 < p_1 < \ldots < p_k = s$ such that $\sigma_N(c_r) = b_{p_{r-1}+1} \ldots b_{p_r}$ for $r = 1, \ldots, k$. Now fix $j \in \{1, \ldots, s\}$ and let r be such that $p_{r-1} < j \leq p_r$. Suppose that $c_r \in \{a_q, \bar{a}_q\}$. Since s_N follows the pattern σ_N, $s_N^{-1}(\{b\}) \cap S^1_q = \{b = y_0 < y_1 < \ldots < y_{p_r - p_{r-1}} = b\}$ and s_N maps $(y_{j-p_{r-1}-1}, y_{j-p_{r-1}})$ homeomorphically onto $\dot{S}^1_\ell$ where $b_j \in \{a_\ell, \bar{a}_\ell\}$. Define t_N on (x_{j-1}, x_j) by $t_N = (s_N|_{(y_{j-p_{r-1}-1}, y_{j-p_{r-1}})})^{-1} \circ f^{n_N}|_{(x_{j-1}, x_j)}$. Extend t_N to a continuous surjection from $[x_{j-1}, x_j]$ onto $[y_{j-p_{r-1}-1}, y_{j-p_{r-1}}]$. Carry out the above for each $i \in \{1, \ldots, n\}$ and $j \in \{1, \ldots, s = s_i\}$. That there is no ambiguity in the definition of t_N is guaranteed by the construction and the fact that $\sigma_N \tau_N = \chi_f^{n_N}$. It is clear that t_N is a continuous strictly piecewise monotone surjection that follows the pattern τ_N and that $s_N t_N = f^{n_N}$. Now define $s_{N+1} : X_m \to X_n$ using t_N and g^{m_N}, etc. The commuting diagram



induces a homeomorphism between $\varprojlim f$ and $\varprojlim g$. □

In order to obtain a converse to Theorem 1.5 we will need to assume that the point $\underline{b} = (b, b, \ldots)$ in $\varprojlim f$ and $\varprojlim g$ is somehow distinguished so that any homeomorphism would necessarily take $\underline{b}$ to $\underline{b}$. We proceed to isolate three ways this can happen for $f, g \in \mathcal{M}$. To this end, consider the action of $f \in \mathcal{M}_{n,n}$ on the "edge germs" at b. There are $2n$ germs at b, $\mathcal{E} = \{e_1, \ldots, e_n, \bar{e}_1, \ldots, \bar{e}_n\}$, where we denote the edge germs of S^1_i at b by e_i and $\bar{e}_i$: e_i (resp. $\bar{e}_i$) emanates from b in the direction of the positive (resp. negative) orientation of S^1_i. Define $f_* : \mathcal{E} \to \mathcal{E}$ by $f_*(e_i) = e_j$ (resp. $\bar{e}_j$) provided $\chi_f(a_i) = a_j \ldots$ (resp. $\bar{a}_j \ldots$) and $f_*(\bar{e}_i) = e_j$ (resp. $\bar{e}_j$) provided $\chi_f(\bar{a}_i) = a_j \ldots$ (resp. $\bar{a}_j \ldots$). Let $\mathcal{R}(f) = \bigcap_{k \geq 0} f_*^k(\mathcal{E})$ be the eventual range of f_*. Note that if $f \in \mathcal{M}_{n,n}$ then $\mathcal{R}(f) = f_*^{2n}(\mathcal{E})$.

By an *open arc* we will mean a homeomorphic copy of the interval $(-1, 1)$.

Lemma 1.6. *If $f \in \mathcal{M}$ and $\#\mathcal{R}(f) = 1$ then there does not exist an open arc in $\varprojlim f$ containing $\underline{b}$.*

Proof. By Lemma 1.4, we may assume that $f \in \mathcal{M}_{n,n}$ is strictly piecewise monotone (the homeomorphisms of Lemmas 1.2 and 1.3 can be chosen to take $\underline{b}$ to $\underline{b}$). Moreover, we may assume that $\#f_*(\mathcal{E}) = 1$, since $\varprojlim f^{2n} \simeq \varprojlim f$, and that $f|_{[b,\epsilon]}$ is the identity for some $\epsilon \neq b$ where $[b, \epsilon]$ is a closed arc in S^1_j and $f_*(\mathcal{E}) = \{e_j\}$ or $\{\bar{e}_j\}$. Then, if α is an open arc in $\varprojlim f$ containing $\underline{b}$, we may assume that $\alpha_k = \pi_k(\alpha) \subset [b, \epsilon]$ for all k. But then $\alpha \subset \varprojlim f|_{[b,\epsilon]}$, and this is clearly not possible since $\underline{b}$ is an endpoint of the closed arc $\varprojlim f|_{[b,\epsilon]}$. □



$\#\mathcal{R}(f) = 1$ | $\#\mathcal{R}(f) = 3$, $\underline{b}$ is a branch point | f folds at b

We will call a point x in a continuum X a *branch point* if there is a continuum $Y \subset X$ such that Y is homeomorphic with the letter Y (three closed arcs joined at an endpoint) and $Y \backslash \{x\}$ has three components.

Lemma 1.7. *If $f \in \mathcal{M}$ and $\#\mathcal{R}(f) \geq 3$ then $\underline{b}$ is a branch point of $\varprojlim f$.*

Proof. Replacing f by $f^{2n!}$, if necessary, we may suppose that there are distinct edge germs $b_i \in \mathcal{E}$ such that $f_*(b_i) = b_i$, for $i = 1, 2, 3$. Say, without loss of generality, that $b_i = e_i$, $i = 1, 2, 3$. We may also assume that $\epsilon_i \in \dot{S}^1_i$ are such that $f|_{[b,\epsilon_i]}$ is the identity for $i = 1, 2, 3$. Let $Y = \varprojlim f|_{\cup_{i=1}^3 [b,\epsilon_i]}$. □

We will say $f \in \mathcal{M}_{n,n}$ *folds at* b if there are $i, j \in \{1, \dots, n\}$ and $k \geq 1$ such that the letters a_j and $\bar{a}_j$ occur consecutively in $\chi_f^k(a_i)$.

Lemma 1.8. *If $f \in \mathcal{M}_{n,n}$ folds at b then no neighborhood of $\underline{b} \in \varprojlim f$ is homeomorphic with the product of a zero-dimensional space and an open arc.*

Proof. We may assume that f_* fixes each edge germ in $\mathcal{R}(f)$ (by replacing f by $f^{2n!}$). For the sake of argument, assume that $f_*(e_1) = e_1$ and that $\bar{a}_1$, a_1 occur consecutively in $\chi_f(a_j)$. Further, assume that f is strictly piecewise monotone and that $f|_{[b,\epsilon]}$ is the identity for some $\epsilon \in \dot{S}^1_1$. Let $a' < c < b' \in \dot{S}^1_j$ be such that $f(c) = b$, $f(a') = f(b') = \epsilon$, $f([a', b']) = [b, \epsilon]$. Let $J_0 = [b, \epsilon]$, $J_1 = [a', b']$, and choose intervals $J_2, J_3, \dots$ such that f takes J_{i+1} homeomorphically onto J_i for $i \geq 1$. Let $X \equiv \{(x_0, x_1, \dots) \in \varprojlim f : \text{for some } n, x_0 = x_1 = \dots = x_n \text{ and } x_{n+k} \in J_k \text{ for } k \geq 1\}$. Then every neighborhood of $\underline{b}$ in $\varprojlim f$ contains a neighborhood of $\underline{b}$ in X, which in turn contains a homeomorphic copy of X. Note that X is homeomorphic with a neighborhood of the point $(0, 1)$ in the "topologist's sine curve" $\{0\} \times [-1, 1] \cup \{(t, sin1/t) : 0 < t \leq 1\}$. No such X is contained in the product of a zero-dimensional set with an arc. □

Following [AM] we will call a closed subset M of a continuum X a *matchbox in* X if there exist a zero-dimensional space S and a topological embedding $h : S \times [-1, 1] \to X$ such that $h(S \times [-1, 1]) = M$ and $h(S \times (-1, 1)) = \dot{M}$, the interior of M. The map $h : S \times [-1, 1] \to M$ is called a *parametrization of* M. An *oriented matchbox* is a matchbox provided with a parametrization, whose matches (arc components) are oriented by the canonical orientation of $[-1, 1]$ via the parametrization. Let $pr_1 : S \times [-1, 1] \to S$ be defined by $pr_1(s, t) = s$ and $pr_2 : S \times [-1, 1] \to [-1, 1]$ be defined by $pr_2(s, t) = t$. If M_1 and M_2 are oriented matchboxes in X with parametrizations $h_1 : S_1 \times [-1, 1] \to M_1$ and $h_2 : S_2 \times [-1, 1] \to M_2$ then we say that M_1 and M_2 are *oriented coherently* (resp. *anticoherently*) if for each $x \in int(M_1 \cap M_2)$ the map $pr_2|_{h_2^{-1}(A_x)} \circ h_2^{-1} \circ h_1 \circ (pr_2|_{h_1^{-1}(A_x)})^{-1} : pr_2 \circ h_1^{-1}(A_x) \to pr_2 \circ h_2^{-1}(A_x)$ is increasing (resp. decreasing), where A_x is the component of $M_1 \cap M_2$ containing x.

Lemma 1.9. *If $f \in \mathcal{M}_{n,n}$ and J is a closed arc of $X_n \backslash \{b\}$ then for $k = 0, 1, \dots$, $\pi_k^{-1}(J)$ is an oriented matchbox in $\varprojlim f \backslash \{\underline{b}\}$.*

Proof. Let $h : [-1,1] \to J$ be an orientation preserving homeomorphism onto J (the orientation of J agreeing with the orientation of the S^1_i containing it). Let $\dot{J}_1, \dots, \dot{J}_\ell$ be the arc components of $X_n \backslash f^{-1}(\{b\})$ and Σ_ℓ be the product space $\prod_{j=1}^{\infty}\{1,\dots,\ell\}$ where $\{1,\dots,\ell\}$ has the discrete topology. Then $S \equiv \{(b_1, b_2, \dots) \in \Sigma_\ell : J \subset f(\dot{J}_{b_1})$ and $\dot{J}_{b_i} \subset f(\dot{J}_{b_{i+1}})$ for $i = 1, 2, \dots\}$ is a closed subset of the Cantor set Σ_ℓ. Let $h_k : S \times [-1,1] \to \pi_k^{-1}(J)$ be the homeomorphism defined by $h_k((b_1, b_2, \dots), t) = (f^k \circ h(t), f^{k-1} \circ h(t), \dots, h(t), (f|_{J_{b_1}})^{-1} \circ h(t), (f|_{J_{b_2}})^{-1} \circ (f|_{J_{b_1}})^{-1} \circ h(t), \dots)$. □

Lemma 1.10. *If $f \in \mathcal{M}$ and $\underline{x} \in \varprojlim f \backslash \{\underline{b}\}$ then there is a matchbox $M \subset \varprojlim f \backslash \{\underline{b}\}$ such that $\underline{x} \in \dot{M}$.*

Proof. Choose $f \in \mathcal{M}_{n,n}$ to be strictly piecewise monotone and let k be such that $\pi_k(\underline{x}) \neq b$. Let J be a closed arc in $X_n \backslash \{b\}$ with $\pi_k(\underline{x}) \in \dot{J}$. Then $M = \pi_k^{-1}(J)$ is a matchbox with $\underline{x} \in \dot{M}$. □

We will say b is *distinguished* for $f \in \mathcal{M}$ if f folds at b or $\#\mathcal{R}(f) \neq 2$. Let $\mathcal{M}^* = \{f \in \mathcal{M} : b \text{ is distinguished for } f\}$.

Proposition 1.11. *If $f, g \in \mathcal{M}^*$ and ψ is a homeomorphism from $\varprojlim f$ to $\varprojlim g$ then $\psi(\underline{b}) = \underline{b}$.*

Proof. This follows immediately from Lemmas 1.6-1.8 and 1.10: $\underline{b}$ is the only point which does not admit a matchbox neighborhood. □

Let $f \in \mathcal{M}_{n,n}$ be such that b is distinguished for f. For each $i = 1, \dots, n$, let I_i be a closed interval in $\dot{S}^1_i$ and $h_i : [-1,1] \to I_i$ be a homeomorphism such that the orientation induced on I_i by the canonical orientation of $[-1,1]$ agrees with the orientation of S^1_i. Let $\chi_f : \mathcal{A}_n \to \mathcal{W}(\mathcal{A}_n)$ be the substitution determined by f. If $\chi_f(a_i) = c_1 \dots c_s$ then for each $j = 0, \dots, s$, we choose $x_{i,j} \in [-1,1]$ such that $x_{i,0} = -1$, $x_{i,s} = 1$, and $x_{i,j-1} < x_{i,j}$ for each $j = 1, \dots, s$. Let $\tilde{f} \in \mathcal{M}_{n,n}$ be such that $\tilde{f}(X_n \backslash \bigcup_{i=1}^n \dot{I}_i) = \{b\}$ and $\tilde{f}$ takes $\dot{I}_{i,j} \equiv h_i((x_{i,j-1}, x_{i,j}))$ homeomorphically onto $\dot{S}^1_\ell$ in an orientation preserving (resp. reversing) way if $c_j = a_\ell$ (resp. $\bar{a}_\ell$). Since $\tilde{f}$ follows the same pattern as f and b is distinguished for $\tilde{f}$, $\varprojlim \tilde{f} \backslash \{\underline{b}\} \simeq \varprojlim f \backslash \{\underline{b}\}$.

Lemma 1.12. *If M is a matchbox in $\varprojlim \tilde{f} \backslash \{\underline{b}\}$ then there exists a K such that $M \subset \bigcup_{i=1}^n \pi_k^{-1}(I_i)$ for every $k \geq K$.*

Proof. For each $\underline{x} \in \varprojlim \tilde{f} \backslash \{\underline{b}\}$, so in particular for each $\underline{x} \in M$, there is a k such that $\pi_j(\underline{x}) \in \bigcup_{i=1}^n \dot{I}_i$ for all $j \geq k$. Then, by compactness of M there is a K such that $\pi_k(\underline{x}) \in \bigcup_{i=1}^n \dot{I}_i$ for all $\underline{x} \in M$ and all $k \geq K$. That is, $M \subset \bigcup_{i=1}^n \pi_k^{-1}(I_i)$ for all $k \geq K$. □

Lemma 1.13. *If M is an oriented matchbox in $\varprojlim \tilde{f} \backslash \{\underline{b}\}$ then there is a positive integer K such that for each $k \geq K$ there is a collection of pairwise disjoint matchboxes $M_1, \dots, M_{\ell_k}$ whose union is M such that, for $j = 1, \dots, \ell_k$, the following conditions hold:*

i) there exists an $i \in \{1, \dots, n\}$ such that $M_j \subset \pi_k^{-1}(I_i)$;
ii) M_j, with the orientation inherited from M, and $\pi_k^{-1}(I_i)$ are oriented coherently or anticoherently;

iii) there is an $\underline{x}_j \in \dot{M}_j$ such that each match of M_j passes through $\pi_k^{-1} \circ \pi_k(\{\underline{x}_j\})$.

Proof. Let $h : S \times [-1,1] \to M$ be a parametrization of M. By Lemma 1.12, there exists an L such that $M \subset \bigcup_{i=1}^n \pi_\ell^{-1}(I_i)$ for all $\ell \geq L$. Given $\underline{x} \in \dot{M}$ and an open set U, where $\underline{x} \in U \subset \dot{M}$, there exists an $\ell \geq L$ such that $\pi_\ell^{-1} \circ \pi_\ell(\{\underline{x}\}) \subset U$. The matchbox $M_{\underline{x},\ell} \equiv h(pr_1 \circ h^{-1} \circ \pi_\ell^{-1} \circ \pi_\ell(\{\underline{x}\}) \times [-1,1]) \subset M$ inherits an orientation from M. For $\underline{z} \in \pi_\ell^{-1} \circ \pi_\ell(\{\underline{x}\})$ and $k \geq \ell$, $M_{\underline{z},k} \subset M_{\underline{z},\ell} = M_{\underline{x},\ell}$. Also for $\underline{w}, \underline{z} \in \pi_\ell^{-1} \circ \pi_\ell(\{\underline{x}\})$ and $k \geq \ell$, $M_{\underline{w},k}$ and $M_{\underline{z},k}$ are either disjoint or equal. So given $k \geq \ell$, there exist $\underline{z}_1, \dots, \underline{z}_{r_k} \in \pi_\ell^{-1} \circ \pi_\ell(\{\underline{x}\})$ such that $M_{\underline{z}_1,k}, \dots, M_{\underline{z}_{r_k},k}$ is a collection of pairwise disjoint matchboxes whose union is $M_{\underline{x},\ell}$. For each $\underline{x} \in C \equiv h(S \times \{0\})$ (cross-section of M), choose $\ell(\underline{x}) \geq L$ such that $\pi_{\ell(\underline{x})}^{-1} \circ \pi_{\ell(\underline{x})}(\{\underline{x}\}) \subset \dot{M}$. So there exist $\underline{x}_1, \dots, \underline{x}_s \in C$ such that $M_{\underline{x}_1,\ell(\underline{x}_1)}, \dots, M_{\underline{x}_s,\ell(\underline{x}_s)}$ is a minimal cover of M (i.e., $M_{\underline{x}_i,\ell(\underline{x}_i)} \not\subset \bigcup_{j \neq i} M_{\underline{x}_j,\ell(\underline{x}_j)}$ for each $i = 1, \dots, s$). Let M_i be the matchbox $M_{\underline{x}_i,\ell(\underline{x}_i)} \setminus \bigcup_{j<i} M_{\underline{x}_j,\ell(\underline{x}_j)}$ and $C_i = \{\underline{z} \in \pi_{\ell(\underline{x}_i)}^{-1} \circ \pi_{\ell(\underline{x}_i)}(\{\underline{x}_i\}) : \underline{z} \in M_i\}$. Choose $k_i \geq \ell(\underline{x}_i)$ such that, for all $\underline{z} \in C_i$, $M_{\underline{z},k_i} \subset M_i$ and $M_{\underline{z},k_i}$ and $\pi_{k_i}^{-1}(I_j)$ are oriented coherently or anticoherently where j is such that I_j contains $\pi_{k_i}(\underline{z})$. Let $k \geq K \equiv \max\{k_1, \dots, k_s\}$. Then there exist $\underline{z}_1, \dots, \underline{z}_{r_k} \in \bigcup_{i=1}^s C_i$ such that $M_{\underline{z}_1,k}, \dots, M_{\underline{z}_{r_k},k}$ is a collection of pairwise disjoint matchboxes whose union is M. □

Lemma 1.14. *If M is a matchbox in $\varprojlim \tilde{f} \setminus \{\underline{b}\}$ that intersects each arc component of $\varprojlim \tilde{f} \setminus \{\underline{b}\}$ then there is a K such that $\pi_k^{-1}(I_i) \cap M \neq \emptyset$ for $k \geq K$ and $i = 1, \dots, n$.*

Proof. Choose L such that $\pi_\ell^{-1} \circ \pi_\ell(\{\underline{x}\}) \subset M \subset \bigcup_{i=1}^n \pi_\ell^{-1}(I_i)$ for each $\underline{x}$ in a cross-section of M and $\ell \geq L$. For purposes of contradiction, assume there exist $L \leq k_1 < k_2 < \dots$ and $I_{i_{k_1}}, I_{i_{k_2}}, \dots$ such that $\pi_{k_j}^{-1}(I_{i_{k_j}}) \cap M = \emptyset$. If $\pi_k^{-1}(I_i) \cap M = \emptyset$ for some i and $k \geq L$ then $\pi_\ell^{-1}(I_j) \cap M = \emptyset$ for each I_j covered by $\tilde{f}^{k-\ell}(I_i)$ where $L \leq \ell \leq k$. So there exist $i_L, i_{L+1}, \dots$ such that $\pi_j^{-1}(I_{i_j}) \cap M = \emptyset$ and $I_{i_j} \subset \tilde{f}(I_{i_{j+1}})$ for $j = L, L+1, \dots$. There exists an $\underline{x} \in \varprojlim \tilde{f} \setminus \{\underline{b}\}$ such that $\pi_j(\underline{x}) \in I_{i_j}$ for $j = L, L+1, \dots$. Choose $\underline{z} \in M$ so that $\underline{x}$ and $\underline{z}$ lie on the same arc component of $\varprojlim \tilde{f} \setminus \{\underline{b}\}$ and let A be the arc between $\underline{x}$ and $\underline{z}$. There is then an $\ell \geq L$ such that $b \notin \pi_\ell(A)$. Thus $\pi_{\ell+1}(A) \subset I_{i_{\ell+1}}$. But this contradicts $\pi_{\ell+1}^{-1}(I_{i_{\ell+1}}) \cap M = \emptyset$. □

Let $g \in \mathcal{M}_{m,m}$ and define the map $\tilde{g} : X_m \to X_m$ as above.

Lemma 1.15. *If $\varprojlim \tilde{f} \setminus \{\underline{b}\} \simeq \varprojlim \tilde{g} \setminus \{\underline{b}\}$ then $\chi_{\tilde{f}} \sim_w \chi_{\tilde{g}}$.*

Proof. Let Ψ be a homeomorphism from $\varprojlim \tilde{f} \setminus \{\underline{b}\}$ to $\varprojlim \tilde{g} \setminus \{\underline{b}\}$. We inductively choose integers $0 = s_1 < t_1 < s_2 < t_2 < \dots$ in the following way. For a given s_k, $\pi_{s_k}^{-1}(I_1), \dots, \pi_{s_k}^{-1}(I_n)$ are pairwise disjoint matchboxes in $\varprojlim \tilde{f} \setminus \{\underline{b}\}$, with orientation given by Lemma 1.9, and $\bigcup_{\ell=1}^n \pi_{s_k}^{-1}(I_\ell)$ intersects each arc component of $\varprojlim \tilde{f} \setminus \{\underline{b}\}$. So $\Psi(\pi_{s_k}^{-1}(I_1)), \dots, \Psi(\pi_{s_k}^{-1}(I_n))$ are pairwise disjoint oriented matchboxes in $\varprojlim \tilde{g} \setminus \{\underline{b}\}$ and $\Psi(\bigcup_{\ell=1}^n \pi_{s_k}^{-1}(I_\ell)) = \bigcup_{\ell=1}^n \Psi(\pi_{s_k}^{-1}(I_\ell))$ intersects each arc component of $\varprojlim \tilde{g} \setminus \{\underline{b}\}$. By Lemmas 1.13 and 1.14, we can choose a $t_k > s_k$ such that, for each $i = 1, \dots, m$, $\pi_{t_k}^{-1}(I_i') \cap \Psi(\bigcup_{\ell=1}^n \pi_{s_k}^{-1}(I_\ell)) \neq \emptyset$ and, for each $j = 1, \dots, n$, $\Psi(\pi_{s_k}^{-1}(I_j))$ is the union of a finite collection of pairwise disjoint matchboxes satisfying conditions $i) - iii)$ of Lemma 1.13. Similarly, for a given t_k, $\Psi^{-1}(\pi_{t_k}^{-1}(I_1')), \dots, \Psi^{-1}(\pi_{t_k}^{-1}(I_m'))$ are pairwise disjoint oriented matchboxes in

$\varprojlim \tilde{f}\backslash\{\underline{b}\}$ and $\Psi^{-1}(\bigcup_{\ell=1}^{m}\pi_{t_k}^{-1}(I'_\ell))$ intersects each arc component of $\varprojlim \tilde{f}\backslash\{\underline{b}\}$. By Lemmas 1.13 and 1.14, we can choose an $s_{k+1} > t_k$ such that, for $j = 1,\dots,n$, $\pi_{s_{k+1}}^{-1}(I_j) \cap \Psi^{-1}(\bigcup_{\ell=1}^{m}\pi_{t_k}^{-1}(I'_\ell)) \neq \emptyset$ and, for each $i = 1,\dots,m$, $\Psi^{-1}(\pi_{t_k}^{-1}(I'_i))$ is the union of a finite collection of pairwise disjoint matchboxes satisfying conditions $i)-iii)$ of Lemma 1.13. For $k = 1, 2\dots$, we define a substitution $\sigma_k : \mathcal{A}_m \to \mathcal{W}(\mathcal{A}_n)$ in the following way. Choose a match A'_i of $\pi_{t_k}^{-1}(I'_i)$. Let $A'_{i,1} < \dots < A'_{i,s}$ be the arc components of $A'_i \cap \Psi(\bigcup_{\ell=1}^{n}\pi_{s_k}^{-1}(I_\ell)) \neq \emptyset$, with the ordering given by the positive orientation of $\pi_{t_k}^{-1}(I'_i)$. Let $\sigma_k(a'_i) = c_1 \dots c_s$ where $c_j = a_\ell$ (resp. $\bar{a}_\ell$) if $\Psi^{-1}(A'_{i,j}) \subset \pi_{s_k}^{-1}(I_\ell)$ and $\Psi^{-1}(A'_{i,j})$ and $\pi_{s_k}^{-1}(I_\ell)$ are oriented coherently (resp. anticoherently). Let $\sigma_k(\bar{a}'_i) = \overline{\sigma_k(a'_i)}$. Conditions $i) - iii)$ of Lemma 1.13 ensure that this substitution is independent of the chosen match. We define $\tau_k : \mathcal{A}_n \to \mathcal{W}(\mathcal{A}_m)$ similarly. Let $n_k = s_{k+1} - s_k$ and $m_k = t_{k+1} - t_k$. If A is a match of $\bigcup_{\ell=1}^{n}\pi_{s_{k+1}}^{-1}(I_\ell)$, then the substitution $\sigma_k \circ \tau_k : \mathcal{A}_n \to \mathcal{W}(\mathcal{A}_n)$ describes the way in which $\Psi^{-1} \circ \Psi(A)$ passes through $\bigcup_{\ell=1}^{n}\pi_{s_k}^{-1}(I_\ell)$, but $(\chi_{\tilde{f}})^{n_k} = (\chi_{\tilde{f}})^{s_{k+1}-s_k} : \mathcal{A}_n \to \mathcal{W}(\mathcal{A}_n)$ also describes the way in which A passes through $\bigcup_{\ell=1}^{n}\pi_{s_k}^{-1}(I_\ell)$. So $\sigma_k \circ \tau_k = (\chi_{\tilde{f}})^{n_k}$. Similarly $\tau_k \circ \sigma_{k+1} = (\chi_{\tilde{g}})^{m_k}$. Thus $\chi_{\tilde{f}} \sim_w \chi_{\tilde{g}}$. □

Theorem 1.16. *Given $f, g \in \mathcal{M}^*$, $\varprojlim f \simeq \varprojlim g$ iff $\chi_f \sim_w \chi_g$.*

Proof. By Theorem 1.5, if $\chi_f \sim_w \chi_g$ then $\varprojlim f \simeq \varprojlim g$. Conversely, if $f, g \in \mathcal{M}^*$ and $\varprojlim f \simeq \varprojlim g$, then $\varprojlim f\backslash\{\underline{b}\} \simeq \varprojlim g\backslash\{\underline{b}\}$ by Proposition 1.11. Let $\tilde{f}$, $\tilde{g}$ be as above. Then $\varprojlim \tilde{f}\backslash\{\underline{b}\} \simeq \varprojlim f\backslash\{\underline{b}\}$ and $\varprojlim g\backslash\{\underline{b}\} \simeq \varprojlim \tilde{g}\backslash\{\underline{b}\}$. So by Lemma 1.15, $\chi_f = \chi_{\tilde{f}} \sim_w \chi_{\tilde{g}} = \chi_g$. □

Let $\mathcal{M}^I = \mathcal{M}\backslash\mathcal{M}^*$. The I in the notation stands for immersion: X_n can be given the structure of a smooth branched manifold and for each $f \in \mathcal{M}^I$ there is a smooth immersion $g \in \mathcal{M}^I$ with $\chi_f = \chi_g$. But this will not concern us here (see [W2],[J]). We will say that $f \in \mathcal{M}_n^I \equiv \mathcal{M}_{n,n} \cap \mathcal{M}^I$ is *orientation preserving* provided, with respect to some orientation on the S_i^1, $\chi_f(a_i) \in \mathcal{W}(\{a_1,\dots,a_n\})$ for each $i \in \{1,\dots,n\}$. Let $\mathcal{M}_o^I = \{f \in \mathcal{M}^I : f^2 \text{ is orientation preserving}\}$. The following theorem is proved in [J].

Theorem 1.17. *If $f, g \in \mathcal{M}_o^I$ and $\varprojlim f \simeq \varprojlim g$ then $A_f \sim_w A_g$.*

It is also proved in [J] that for $f \in \mathcal{M}^I\backslash\mathcal{M}_o^I$, say $f : X_n \to X_n$, there is an associated $f' \in \mathcal{M}_o^I$, $f' : X_m \to X_m$. Moreover, if $\varprojlim f \simeq \varprojlim g$ for $f, g \in \mathcal{M}^I\backslash\mathcal{M}_o^I$, then $\varprojlim f' \simeq \varprojlim g'$, which with Theorem 1.17, yields the following.

Theorem 1.18. *If $f, g \in \mathcal{M}^I\backslash\mathcal{M}_o^I$ and $\varprojlim f \simeq \varprojlim g$ then $A_{f'} \sim_w A_{g'}$.*

A square matrix A with nonnegative entries is *aperiodic* provided $A^k > 0$ for some k. Such a matrix has a simple real eigenvalue λ_A, greater in modulus than all other eigenvalues, called the *Perron eigenvalue* of A. Let $\mathbb{Q}(\lambda_A)$ denote the extension field of the rationals $\mathbb{Q}$ by λ_A.

Lemma 1.19. *If A, B are aperiodic square nonnegative integral matrices and $A \sim_w B$ then $\mathbb{Q}(\lambda_A) = \mathbb{Q}(\lambda_B)$.*

Proof. See [BD]. □

Corollary 1.20. *Let $f, g \in \mathcal{M}$ and suppose that A_f and A_g are aperiodic. If $\varprojlim f \simeq \varprojlim g$ then $\mathbb{Q}(\lambda_A) = \mathbb{Q}(\lambda_B)$.*

Proof. If $f, g \in \mathcal{M}^*$, this follows from Lemmas 1.1, 1.19 and Theorem 1.16. If $f, g \in \mathcal{M}_o^I$, this follows from Theorem 1.17 and Lemma 1.19. If $f, g \in \mathcal{M}^I \backslash \mathcal{M}_o^I$, Jacklitch [J] proves that $\lambda_{f'} = \lambda_f$, $\lambda_{g'} = \lambda_g$, so the corollary follows from Theorem 1.18 and Lemma 1.19. There are no other possibilities for $\varprojlim f \simeq \varprojlim g$. □

The proofs of Theorems 1.17 and 1.18 are quite different. Nonetheless, we make the following conjecture.

Conjecture 1.21. *Given $f, g \in \mathcal{M}$, $\varprojlim f \simeq \varprojlim g$ iff $\chi_f \sim_w \chi_g$.*

2. Applications

In this section we present several applications of the preceding theorems. The first of these will be to one-dimensional substitution tilings and for this we borrow heavily from [AP].

Consider a substitution $\chi : \mathcal{A} = \{a_1, \dots, a_n\} \to \mathcal{W}(\mathcal{A})$, $n \geq 2$, with associated matrix A_χ. To avoid degeneracy, we assume that for no word w and positive integer k is $\chi^k(w) = ww \dots w$. We also assume that A_χ is aperiodic with Perron eigenvalue λ_χ and associated positive (left) eigenvector $\vec{v}_\chi$ with entries $\lambda_1, \dots, \lambda_n$. The intervals $P_i = [0, \lambda_i]$, $i = 1, \dots, n$, are called prototiles (consider P_i to be distinct from P_j for $i \neq j$ even if $\lambda_i = \lambda_j$). A *tiling* T of $\mathbb{R}$ by the prototiles is a collection $T = \{T_i\}_{i=-\infty}^{\infty}$ of tiles T_i, $\bigcup_{i=-\infty}^{\infty} T_i = \mathbb{R}$, with each T_i a translate of some P_j and with $T_i \cap T_{i+1}$ a singleton for each i.

Now $\lambda_\chi \lambda_i = \sum_{j=1}^{r} \lambda_{i(j)}$ where $i(j) = k$ provided $\chi(a_i) = c_1 \dots c_r$ with $c_j = a_k$. Thus $\lambda_\chi P_i = \bigcup_{s=0}^{r-1} (\sum_{j=1}^{s} \lambda_{i(j)}) + P_{i(s+1)}$ that is, $\lambda_\chi P_i$ is tiled by $\{T_s\}_{s=0}^{r-1}$, $T_s = (\sum_{j=1}^{s} \lambda_{i(j)}) + P_{i(s+1)}$. This process is called inflation and substitution and extends to a map σ_χ taking a tiling $T = \{T_i\}_{i=-\infty}^{\infty}$ of $\mathbb{R}$ by prototiles to a new tiling, $\sigma_\chi(T)$ of $\mathbb{R}$ by prototiles defined by inflating, substituting, and suitably translating each T_i.

There is a natural topology on the collection Σ_χ of all tilings of $\mathbb{R}$ by prototiles ($\{T_i\}_{i=-\infty}^{\infty}$ and $\{T_i'\}_{i=-\infty}^{\infty}$ are "close" if there is an ϵ near zero and a large N so that if $0 \in T_k \cap T_\ell'$ then $T_{k+i} = \epsilon + T_{\ell+i}'$ for $i = -N, \dots, N$, see [AP] for details). Σ_χ is a compact metrizable space with this topology and $\sigma_\chi : \Sigma_\chi \to \Sigma_\chi$ is continuous. Let $T \in \Sigma_\chi$ be any tiling that is periodic under σ_χ (there are infinitely many such) and consider the orbit, $\mathcal{O}(T)$, of T under the flow $\phi_t : \Sigma_\chi \to \Sigma_\chi$ by $\phi_t(\{T_i\}_{i=-\infty}^{\infty}) = \{t + T_i\}_{i=-\infty}^{\infty}$. Let $\mathcal{T}_\chi = cl(\mathcal{O}(T))$; $\mathcal{T}_\chi$ is the *tiling space associated with* χ. Then $\mathcal{T}_\chi$ does not depend on the T chosen above, ϕ_t is minimal on $\mathcal{T}_\chi$ (so in particular, $\mathcal{T}_\chi$ is a continuum) and $\sigma_\chi : \mathcal{T}_\chi \to \mathcal{T}_\chi$ is a homeomorphism.

Given two such substitutions χ, ψ, we wish to apply theorems from the previous section in an effort to distinguish $\mathcal{T}_\chi$ from $\mathcal{T}_\psi$ if they are nonhomeomorphic. In order to (partially) accomplish this, we need to consider *collared* matrices $A_{\tilde{\chi}}$ and $A_{\tilde{\psi}}$ associated with χ and ψ. Consider the collection $\tilde{\mathcal{A}} = \tilde{\mathcal{A}}_\chi = \{(a_i a_j a_k) : a_i a_j a_k \text{ occurs as a subword of } \chi^\ell(a_s), \text{ some } \ell, s\}$. Define $\tilde{\chi} : \tilde{\mathcal{A}} \to \mathcal{W}(\tilde{\mathcal{A}})$ by $\tilde{\chi}((a_i a_j a_k)) = (b_0 b_1 b_2)(b_1 b_2 b_3) \dots (b_{s-2} b_{s-1} b_s)(b_{s-1} b_s b_{s+1})$ where $\chi(a_i) = \dots b_0$, $\chi(a_j) = b_1 \dots b_s$, and $\chi(a_k) = b_{s+1} \dots$. Let $A_{\tilde{\chi}}$ be the matrix associated with $\tilde{\chi}$ (choose any numbering of the elements of $\tilde{\mathcal{A}}$). Then $A_{\tilde{\chi}}$ is also aperiodic. Let $\lambda_{\tilde{\chi}}$ be the Perron eigenvalue of $A_{\tilde{\chi}}$.

Lemma 2.1. $\lambda_{\tilde{\chi}} = \lambda_\chi$.

Proof. By construction of $\tilde{\chi}$, there is a nonnegative integer matrix P of rank n such that $PA_{\tilde{\chi}} = A_\chi P$. Let $\tilde{v}$ be a positive eigenvector of $A_{\tilde{\chi}}$ corresponding to the eigenvalue $\lambda_{\tilde{\chi}}$. Then $A_\chi P\tilde{v} = PA_{\tilde{\chi}}\tilde{v} = P\lambda_{\tilde{\chi}}\tilde{v} = \lambda_{\tilde{\chi}}P\tilde{v}$. Thus $v = P\tilde{v}$ is a positive eigenvector for A_χ with eigenvalue $\lambda_{\tilde{\chi}}$. Thus $\lambda_{\tilde{\chi}} = \lambda_\chi$ by the Perron-Frobenius Theorem. □

Theorem 2.2. *If $\mathcal{T}_\chi \simeq \mathcal{T}_\psi$ then $A_{\tilde{\chi}} \sim_w A_{\tilde{\psi}}$.*

Proof. Anderson and Putnam ([AP]) prove that there are branched one-manifolds G_χ and G_ψ and expanding, orientation preserving immersions $f_\chi : G_\chi \to G_\chi$ and $f_\psi : G_\psi \to G_\psi$ such that $\mathcal{T}_\chi \simeq \varprojlim f_\chi$ and $\mathcal{T}_\psi \simeq \varprojlim f_\psi$. Furthermore, the Markov matrices associated with f_χ and f_ψ are $A_{\tilde{\chi}}$ and $A_{\tilde{\psi}}$, respectively. Using the Williams moves to split the branched manifolds into wedges of circles, there are $g_\chi : X_n \to X_n$ and $g_\psi : X_m \to X_m$ with $\varprojlim g_\chi \simeq \varprojlim f_\chi$ and $\varprojlim g_\psi \simeq \varprojlim f_\psi$ (see [W2] and [J]). Moreover, $g_\chi, g_\psi \in \mathcal{M}^I_{o.p.}$ and the associated matrices A_{g_χ} and A_{g_ψ} are shift equivalent to some powers of $A_{\tilde{\chi}}$ and $A_{\tilde{\psi}}$, respectively. In particular $A_{g_\chi} \sim_w A_{\tilde{\chi}}$ and $A_{g_\psi} \sim_w A_{\tilde{\psi}}$. Assuming $\mathcal{T}_\chi \simeq \mathcal{T}_\psi$, it follows from Theorem 1.17 that $A_{\tilde{\chi}} \sim_w A_{\tilde{\psi}}$. □

Corollary 2.3. *If $\mathcal{T}_\chi \simeq \mathcal{T}_\psi$ then $\mathbb{Q}(\lambda_\chi) = \mathbb{Q}(\lambda_\psi)$.*

Proof. This follows from Lemmas 1.19, 2.1 and Theorem 2.2. □

Restricting Conjecture 1.21 to this setting we have

Conjecture 2.4. *$\mathcal{T}_\chi \simeq \mathcal{T}_\psi$ iff $\tilde{\chi} \sim_w \tilde{\psi}$.*

The next application we consider is to one-dimensional unstable manifolds of hyperbolic basic sets. We will assume that combinatorial descriptions of the unstable manifolds are given in terms of collections of Markov rectangles satisfying certain felicitous properties. This situation arises, for example, in [BH] and [BL]. Perhaps the properties R1)-R4) below can always be arranged for the right choice of rectangles in the presence of a local product structure ([HK]).

Suppose that $F : M \to M$ is a diffeomorphism of the k-dimensional manifold M with zero-dimensional hyperbolic invariant set Λ_F having one-dimensional unstable manifold $W^u(\Lambda_F)$. In addition, we suppose that there are pairwise disjoint *rectangles* $R_1, \ldots, R_n \subset M$ with $\Lambda_F \subset \bigcup_{i=1}^n \dot{R}_i$ having the following properties.

R1) Each R_i is parametrized by a homeomorphism $\phi_i : [-1,1] \times [-1,1]^{k-1} \to R_i$.
R2) $\Lambda_F = \bigcap_{\ell=-\infty}^{\infty} F^\ell(\bigcup_{i=1}^n R_i)$ and $R_i \cap \Lambda_F \neq \emptyset$ for each i.
R3) If $z = \phi_i(t,x) \in R_i$ and $F(z) = \phi_j(t',x') \in R_j$, then $F(\phi_i(\{t\} \times [-1,1]^{k-1})) \subset \phi_j(\{t\}' \times [-1,1]^{k-1})$.
R4) For $z \in \Lambda_F$, let $W^u(z,R_i)$ be the component of $W^u(z) \cap R_i$ containing z. Then $W^u(z,R_i)$ meets each $\phi_i(\{t\} \times [-1,1]^{k-1})$ exactly once and if $F(z) \in R_j$ then $F(W^u(z,R_i)) \supset W^u(F(z),R_j)$.
R5) If $z \in (\bigcup_{i=1}^n R_i) \cap (F^{-1}(M \backslash (\bigcup_{i=1}^n R_i))$ then $F^\ell(z) \in M \backslash (\bigcup_{i=1}^n R_i)$ for all $\ell \geq 1$.

Given $(M, F, \Lambda_F, \{R_i\}_{i=1}^n)$ satisfying R1)-R5), let $P_i : R_i \to L_i = [-1,1] \times \{i\}$ be defined by $P_i(\phi_i(t,x)) = (t,i)$. For each $i \in \{1, \ldots, n\}$, let $J_1^i < J_2^i < \ldots < J_{k_i}^i$ be the components of $P_i(R_i \cap F^{-1}(\bigcup_{j=1}^n R_j))$ (ordered according to the usual order on $[-1,1]$). Let $f : \bigcup_{i=1}^n (\bigcup_{j=1}^{k_i} J_j^i) \to \bigcup_{i=1}^n L_i$ be defined by $f((t,i)) = P_j(F(z))$

where $P_i(z) = (t, i)$ and $F(z) \in R_j$. Then f is a continuous surjection. Moreover, for each $j \in \{1, \dots, k_i\}$ and $i \in \{1, \dots, n\}$ there is an L_ℓ such that $f|_{J_j^i} : J_j^i \to L_\ell$ is a continuous monotone surjection. We define the substitution $\chi_F : \mathcal{A}_n \to \mathcal{W}(\mathcal{A}_n)$ by $\chi_F(a_i) = b_1 \dots b_{k_i}$ with $b_j = a_\ell$ if $f(J_j^i) = L_\ell$ and f is orientation preserving on J_j^i and $b_j = \bar{a}_\ell$ if $f(J_j^i) = L_\ell$ and f is orientation reversing on J_j^i.

Proposition 2.5. *If $(M, F, \Lambda_F, \{R_i\}_{i=1}^n)$ satisfies R1)-R5) and $f' \in \mathcal{M}_{n,n}$ follows the pattern of χ_F then $\varprojlim f' \backslash \{\underline{b}\} \simeq W^u(\Lambda_F)$.*

Proof. Let $h_i : L_i \to I_i \subset \dot{S}_i^1 \subset X_n$ be orientation preserving homeomorphisms for $i = 1, \dots, n$. Let $f' : \bigcup_{i=1}^n h_i(\bigcup_{j=1}^{k_i} J_j^i) \to \bigcup_{i=1}^n I_i$ by $f'(t) = h_j \circ f \circ h_i^{-1}(t)$, i and j as appropriate. Extend f' to a map in $\mathcal{M}_{n,n}$ that follows the pattern of χ_F and so that for $x \notin \mathcal{I} \equiv \bigcup_{i=1}^n I_i$ there is an ℓ such that $(f')^\ell(x) = b$. Let $\mathcal{R} = \bigcup_{i=1}^n R_i$ and $P : \mathcal{R} \to \mathcal{I}$ be defined by $P(z) = h_i \circ P_i(z)$. Then the diagram

$$\begin{array}{ccc} F^{-1}(\mathcal{R}) \cap \mathcal{R} & \xrightarrow{F} & \mathcal{R} \\ \Big\downarrow P & & \Big\downarrow P \\ (f')^{-1}(\mathcal{I}) \cap \mathcal{I} & \xrightarrow[f']{} & \mathcal{I} \end{array}$$

commutes. Given $z \in W^u(\Lambda_F, \mathcal{R}) \equiv \bigcup_{i=1}^n (\bigcup_{z \in \Lambda_F \cap R_i} (W^u(z, R_i)))$, let $H(z) = (P(z), P(F^{-1}(z)), P(F^{-2}(z)), \dots)$. Then $H(z) \in \varprojlim f'$ and $\pi_\ell(H(z)) \in \mathcal{I}$ for $\ell = 0, 1, \dots$.

Claim: $H : W^u(\Lambda_F, \mathcal{R}) \to \pi_0^{-1}(\mathcal{I})$ is a homeomorphism.

Proof: Suppose that $H(z) = H(w)$ for some $z, w \in W^u(\Lambda_F, \mathcal{R})$. Then $P(F^{-\ell}(z)) = P(F^{-\ell}(w))$ for all ℓ. Then $F^{-t}(z), F^{-t}(w) \in \bigcap_{\ell \geq 0} F^\ell(P^{-1}(P(F^{-t-\ell}(\{z\}))))$ for each $t = 0, 1, \dots$. Given $\epsilon > 0$ there is an N such that $\operatorname{diam}(F^N(W^s(v, R_i))) < \epsilon$ for all $v \in \Lambda_F \cap R_i$ and $i = 1, \dots, n$. So there is a $\delta > 0$ such that if the Hausdorff distance between $P^{-1}(P(F^{-N}(\{z\})))$ and $W^s(v, R_i)$ is less than δ then $\operatorname{diam}(F^N(P^{-1}(P(F^{-N}(\{z\}))))) < \epsilon$. Since there are $z', w' \in \Lambda_F$ with $z \in W^u(z')$ and $w \in W^u(w')$, for sufficiently large ℓ, the Hausdorff distance between $P^{-1}(P(F^{-\ell}(\{z\})))$ and $W^s(F^{-\ell}(z'), R_{i_\ell})$ and between $P^{-1}(P(F^{-\ell}(\{w\})))$ and $W^s(F^{-\ell}(w'), R_{i_\ell})$ is less than δ (i_ℓ such that $F^{-\ell}(z'), F^{-\ell}(w') \in R_{i_\ell}$). It follows that the distance between $F^{-t}(z)$ and $F^{-t}(w)$ goes to zero as $t \to \infty$. Thus $z' = w'$. Since $P(z) = P(w)$ and $W^u(z', R_i)$ hits each $\{t\} \times [-1, 1]^{k-1} = P_i^{-1}(\{t\})$ in a unique point, $z = w$. H is surjective because f' follows the pattern $\chi_F : \pi_0^{-1}(\mathcal{I}) = \{\underline{x} \in \varprojlim f' : \pi_k(\underline{x}) \in \mathcal{I}\ \forall k\}$ so given $\underline{x} \in \pi_0^{-1}(\mathcal{I})$ pick $z \in \bigcap_{\ell \geq 0} F^{-\ell}(P^{-1}(\{x_\ell\}))$; then $H(z) = \underline{x}$. Note that for $z \in H^{-1}(W^u(\Lambda_F, \mathcal{R}))$, $H \circ F(z) = \hat{f}'(H(z))$. Extend H to all of $W^u(\Lambda_F)$ by conjugation; that is, for $z \in W^u(\Lambda_F)$ let ℓ be such that $F^{-\ell}(z) \in W^u(\Lambda_F, \mathcal{R})$. Put $H(z) = (\hat{f}')^\ell(H(F^{-\ell}(z))$. Then (by the above note) H is well defined on $W^u(\Lambda_F)$ and, by assumption R5), H is continuous. Clearly $H(W^u(\Lambda_F)) = \bigcup_{\ell \geq 0} (\hat{f}')^\ell(\pi_0^{-1}(\mathcal{I})) = \varprojlim f' \backslash \{\underline{b}\}$. If $\{\underline{z}^\ell\}_{\ell=1}^\infty$ is a sequence in $W^u(\Lambda_F)$ without a limit point in $W^u(\Lambda_F)$ then the (minimal) j_ℓ for which $\underline{z}^\ell \in F^{j_\ell}(W^u(\Lambda_F, \mathcal{R}))$ diverge to ∞. Thus the sequence $H(\underline{z}^\ell) \to \underline{b}$ as $\ell \to \infty$ and it follows H^{-1} is continuous. ∎

With the assumptions of the proposition, we see that the one-point compactification $(W^u(\Lambda_F))^*$ is homeomorphic with $\varprojlim f'$. We will say that $W^u(\Lambda_F)$ is *distinguished at* ∞ if $f' \in \mathcal{M}^*$.

For $z \in \Lambda_F$, an arc component A of $W^u(z)\backslash\{z\}$ is a *free separatrix* of $W^u(\Lambda_F)$ provided $A \cap \Lambda_F = \emptyset$. The reader may check that the number of free separatrices of $W^u(\Lambda_F)$ is precisely $\#\mathcal{R}(f')$, the cardinality of the eventual range of f'_* acting on the edge germs of X_n. Thus we see that, for $(M, F, \Lambda_F, \{R_i\}_{i=1}^n)$ satisfying R1)-R5), $W^u(\Lambda_F)$ is distinguished at ∞ if and only if $W^u(\Lambda_F)$ has other than exactly two free separatrices or, for some ℓ, i, j, $\chi_F^\ell(a_i)$ contains consecutive letters a_j and $\bar{a}_j$ (that is, $F^\ell(R_i)$ passes through R_j then folds back immediately through R_j).

Theorem 2.6. *Suppose that $(M, F, \Lambda_F, \{R_i\}_{i=1}^n)$ and $(N, G, \Lambda_G, \{R'_i\}_{i=1}^m)$ satisfy R1)-R5) and in addition suppose that $W^u(\Lambda_F)$ and $W^u(\Lambda_G)$ are distinguished at ∞ then $W^u(\Lambda_F) \simeq W^u(\Lambda_G)$ if and only if $\chi_F \sim_w \chi_G$.*

Proof. If $W^u(\Lambda_F) \simeq W^u(\Lambda_G)$ then $(W^u(\Lambda_F))^* \simeq (W^u(\Lambda_G))^*$ so that, with f', g' as in Proposition 2.5, $\varprojlim f' \simeq \varprojlim g'$. From Theorem 1.16, $\chi_{f'} \sim_w \chi_{g'}$ so that $\chi_F \sim_w \chi_G$. Conversely, if $\chi_F \sim_w \chi_G$ then $\chi_{f'} \sim_w \chi_{g'}$ so that $\varprojlim f' \simeq \varprojlim g'$, by Theorem 1.16. Since $f', g' \in \mathcal{M}^*$, a homeomorphism of $\varprojlim f'$ to $\varprojlim g'$ must take $\underline{b}$ to $\underline{b}$ (Proposition 1.11). Thus $W^u(\Lambda_F) \simeq \varprojlim f'\backslash\{\underline{b}\} \simeq \varprojlim g'\backslash\{\underline{b}\} \simeq W^u(\Lambda_G)$. □

The idea of studying one-dimensional unstable manifolds by means of their underlying substitutions comes from [V], in which Vago gives a complete and easy-to-compute classification of the unstable manifolds of one-rectangle systems on surfaces, up to homeomorphism and up to conjugacy of the underlying dynamics.

As a final application, consider the *tent maps* $T_s : [0,1] \to [0,1]$ defined by

$$T_s(x) = \begin{cases} sx + 2 - s & , \quad 0 \le x \le \frac{s-1}{s} \\ -sx + s & , \quad \frac{s-1}{s} \le x \le 1 \end{cases} \text{, for } 1 < s \le 2.$$

Ingram has conjectured that $\varprojlim T_{s_1} \not\simeq \varprojlim T_{s_2}$ for $1 < s_1 < s_2 \le 2$.



Consider the set $\mathcal{P} \subset (1,2]$ of parameters for which the critical point $c_s = \frac{s-1}{s}$ is periodic under T_s. Say $s \in \mathcal{P}$ and c_s has period $n+1$ under T_s. Then $\mathcal{O}(c_s) \equiv \{0 = x_0 < \ldots < x_n = 1\}$ breaks $[0,1]$ into n subintervals $J_i = [x_{i-1}, x_i]$, $i = 1, \ldots, n$. For such an s, let $\chi_s : \mathcal{A}_n \to \mathcal{W}(\mathcal{A}_n)$ be the substitution defined by: $\chi_s(a_i) = a_\ell a_{\ell+1} \ldots a_{\ell+k}$ if $T_s(J_i) = J_\ell \cup \ldots \cup J_{\ell+k}$ and T_s is increasing on J_i; $\chi_s(a_i) = \bar{a}_{\ell+k} \ldots \bar{a}_\ell$ if $T_s(J_i) = J_\ell \cup \ldots \cup J_{\ell+k}$ and T_s is decreasing on J_i.

Theorem 2.7. *Given $s_1, s_2 \in \mathcal{P}$, $\varprojlim T_{s_1} \simeq \varprojlim T_{s_2}$ iff $\chi_{s_1} \sim_w \chi_{s_2}$.*

Proof. Given $s \in \mathcal{P}$ with c_s of period $n+1$ under T_s, let $P_s : [0,1] \to X_n = \bigvee_{i=1}^n S_i^1$ take $\dot{J}_i$ (see notation introduced before this theorem) homeomorphically, and in an orientation preserving manner, onto $\dot{S}_i^1$ for each i, and let $P_s(x_i) = b$ for $i = 0, \ldots, n$. Then there is an $f_s \in \mathcal{M}_{n,n}$ such that $P_s \circ T_s = f_s \circ P_s$.

Moreover, $\chi_s = \chi_{f_s}$. Let $\underline{c}_s$ denote the point in $\varprojlim T_s$ with $\pi_{k(n+1)}(\underline{c}_s) = c_s$ for $k = 0, 1, \dots$ and let $\mathcal{O}(\underline{c}_s) \equiv \{\hat{T}^k(\underline{c}_s) : k = 0, \dots, n\}$ be its orbit. Then the map $\hat{P}_s : \varprojlim T_s \to \varprojlim f_s$ defined by $\hat{P}_s((x_0, x_1, \dots)) = (P_s(x_0), P_s(x_1), \dots)$ is a surjection and is one-to-one except on $\mathcal{O}(\underline{c}_s)$ where $\hat{P}(\mathcal{O}(\underline{c}_s)) = \{\underline{b}\}$. For $s \in \mathcal{P}$, the endpoints of the continuum $\varprojlim T_s$ are exactly the points of $\mathcal{O}(\underline{c}_s)$ (see [BM]). For $s_1, s_2 \in \mathcal{P}$, any homeomorphism from $\varprojlim T_{s_1}$ to $\varprojlim T_{s_2}$ must take endpoints to endpoints so such a homeomorphism would descend to a homeomorphism of $\varprojlim f_{s_1}$ with $\varprojlim f_{s_2}$ by way of $\hat{P}_{s_1}$, $\hat{P}_{s_2}$. Moreover, the arc components of the endpoints of $\varprojlim T_s$ are in one-to-one correspondence with elements of $\mathcal{R}(f_s)$, the eventual range of the action of f_s on edge germs. Since c_s is not periodic of period 2 for any $s \in \mathcal{P}$, $\#\mathcal{R}(f_s) \neq 2$ for any $s \in \mathcal{P}$. That is $f_s \in \mathcal{M}^*$ for $s \in \mathcal{P}$. Thus a homeomorphism of $\varprojlim f_{s_1}$ with $\varprojlim f_{s_2}$ would necessarily take $\underline{b}$ to $\underline{b}$ and would then lift to a homeomorphism of $\varprojlim T_{s_1}$ with $\varprojlim T_{s_2}$. Thus for $s_1, s_2 \in \mathcal{P}$, $\varprojlim T_{s_1} \simeq \varprojlim T_{s_2} \Leftrightarrow \varprojlim f_{s_1} \simeq \varprojlim f_{s_2} \Leftrightarrow \chi_{f_{s_1}} \sim_w \chi_{f_{s_2}} \Leftrightarrow \chi_{s_1} \sim_w \chi_{s_2}$. □

The above sort of reasoning can be extended to prove that for s_1, s_2 in the *post-critically-finite* set ($\mathcal{O}(c_s)$ finite, $1 < s \leq 2$), $\varprojlim T_{s_1} \simeq \varprojlim T_{s_2}$ if and only if $\chi_{s_1} \sim_w \chi_{s_2}$. Theorem 2.7 generalizes the result of [BD]: if $s_1, s_2 \in \mathcal{P}$ and $\varprojlim T_{s_1} \simeq \varprojlim T_{s_2}$ then $A_{\chi_{s_1}} \sim_w A_{\chi_{s_2}}$. Bruin [Bru] has recently shown that, for s_1 and s_2 in the post-critically finite set, if $\varprojlim T_{s_1} \simeq \varprojlim T_{s_2}$ then $\log s_1 / \log s_2 \in \mathbb{Q}$.

References

[AM] J.M. Aarts and M. Martens, *Flows on one-dimensional spaces*, Fundamenta Mathematicae, **131** (1988), 53-67.

[AP] J.E. Anderson and I.F. Putnam, *Topological invariants for substitution tilings and their associated C^*-algebras*, Ergod. Th. & Dynam. Sys., **18** (1998), 509-537.

[BD] M. Barge and B. Diamond, *Homeomorphisms of inverse limit spaces of one-dimensional maps*, Fundamenta Mathematicae, **146** (1995), 171-187.

[BH] M. Barge and S. Holte, *Nearly one-dimensional attractors and inverse limits*, Nonlinearity, **8** (1995), 29-42.

[BM] M. Barge and J. Martin, *Endpoints of inverse limit spaces and dynamics*, Continua (Lecture Notes in Pure and Applied Math., 170), Eds Cook et al. Dekker, New York, 1995, 165-182.

[BL] C. Bonatti and R. Langevin, *Difféomorphismes de Smale des surfaces*, Astérisque, **250** (1998).

[BJKR] O. Bratteli, P.E.T. Jørgensen, K.H. Kim and F.W. Roush, *Non-stationarity of isomorphism between AF algebras defined by stationary Bratteli diagrams*, Preprint.

[Bro] M. Brown, *Some applications of an approximation theorem for inverse limits*, Proc. Amer. Math. Soc, **11** (1960), 478-483.

[Bru] H. Bruin, *Inverse limit spaces of post-critically finite tent maps*, Preprint.

[HK] B. Hasselblatt and A. Katok, *Introduction to the Modern Theory of Smooth Dynamical Systems*, Cambridge University Press, Cambridge, New York, 1994.

[J] J. Jacklitch, *Homeomorphisms of one-dimensional hyperbolic attractors*, in preparation.

[SV] R. Swanson and H. Volkmer, *Invariants of weak equivalence in primitive matrices*, Preprint.

[V] G.M. Vago, *Variétés instables d'ensembles hyperboliques*, Thèse, Université de Bourgogne, 1998.

[W1] R.F. Williams, *One-dimensional nonwandering sets*, Topology, **6** (1967), 473-487.

[W2] R.F. Williams, *Classification of one-dimensional attractors*, Proc. Symp. Pure Math., **14** (1970), 341-361.

Department of Mathematical Sciences, Montana State University, Bozeman, Montana 59717
E-mail address: umsfbar@math.montana.edu

E-mail address: jacklitc@math.montana.edu

Laboratoire de Topologie, Département de Mathématiques, Université de Bourgogne, Dijon, France
E-mail address: vago@u-bourgogne.fr